

LINEAR RESPONSE THEORY FOR RANDOM SCHRÖDINGER OPERATORS AND NONCOMMUTATIVE INTEGRATION

NICOLAS DOMBROWSKI AND FRANÇOIS GERMINET

ABSTRACT. We consider an ergodic Schrödinger operator with magnetic field within the non-interacting particle approximation. Justifying the linear response theory, a rigorous derivation of a Kubo formula for the electric conductivity tensor within this context can be found in a recent work of Bouclet, Germinet, Klein and Schenker [BoGKS]. If the Fermi level falls into a region of localization, the well-known Kubo-Středa formula for the quantum Hall conductivity at zero temperature is recovered. In this review we go along the lines of [BoGKS] but make a more systematic use of noncommutative L^p -spaces, leading to a somewhat more transparent proof.

CONTENTS

1. Introduction	1
2. Covariant measurable operators and noncommutative L^p -spaces	3
2.1. Observables	3
2.2. Noncommutative integration	4
2.3. Commutators of measurable covariant operators	5
2.4. Differentiation	6
3. The setting: Schrödinger operators and dynamics	6
3.1. Magnetic Schrödinger operators and time-dependent operators	6
3.2. Adding the randomness	8
4. Linear response theory and Kubo formula	10
4.1. Adiabatic switching of the electric field	10
4.2. The current and the conductivity	12
4.3. Computing the linear response: a Kubo formula for the conductivity	13
4.4. The Kubo-Středa formula for the Hall conductivity	15
References	17

1. INTRODUCTION

In [BoGKS] the authors consider an ergodic Schrödinger operator with magnetic field, and give a controlled derivation of a Kubo formula for the electric conductivity tensor, validating the linear response theory within the noninteracting particle approximation. For an adiabatically switched electric field, they then recover the expected expression for the quantum Hall conductivity whenever the Fermi energy lies either in a region of localization of the reference Hamiltonian or in a gap of the spectrum.

2000 *Mathematics Subject Classification.* Primary 82B44; Secondary 47B80, 60H25.

The aim of this paper is to provide a pedestrian derivation of [BoGKS]’s result and to simplify their “mathematical apparatus” by resorting more systematically to noncommutative L^p -spaces. We also state results for more general time dependent electric fields, so that AC-conductivity is covered as well. That von Neumann algebra and noncommutative integration play an important rôle in the context of random Schrödinger operators involved with the quantum Hall effect goes back to Bellissard, e.g. [B, BES, SB1, SB2].

The electric conductivity tensor is usually expressed in terms of a “Kubo formula,” derived via formal linear response theory. In the context of disordered media, where Anderson localization is expected (or proved), the electric conductivity has driven a lot of interest coming from several perspectives. For time reversal systems and at zero temperature, the vanishing of the direct conductivity is a physically meaningful evidence of a localized regime [FS, AG]. Another direction of interest is the connection between direct conductivity and the quantum Hall effect [ThKNN, St, B, Ku, BES, AvSS, AG]. On the other hand the behaviour of the alternative conductivity at small frequencies within the region of localization is dictated by the celebrated Mott formula [MD] (see [KLP, KLM, KM] for recent important developments). Connected to conductivities, current-current correlations functions have recently drawn a lot of attention as well (see [BH, CGH] and references therein).

During the past two decades a few papers managed to shed some light on these derivations from the mathematical point of view, e.g., [P, Ku, B, NB, AvSS, BES, SB1, SB2, AG, Na, ES, CoJM, CoNP]. While a great amount of attention has been brought to the derivation from a Kubo formula of conductivities (in particular of the quantum Hall conductivity), and to the study of these conductivities, not much has been done concerning a controlled derivation of the linear response and the Kubo formula itself; only the recent papers [SB2, ES, BoGKS, CoJM, CoNP] deal with this question.

In this note, the accent is put on the derivation of the linear response for which we shall present the main elements of proof, along the lines of [BoGKS] but using noncommutative integration. The required material is briefly presented or recalled from [BoGKS]. Further details and extended proofs will be found in [Do]. We start by describing the noncommutative L^p -spaces that are relevant in our context, and we state the properties that we shall need (Section 2). In Section 3 we define magnetic random Schrödinger operators and perturbations of these by time dependent electric fields, but in a suitable gauge where the electric field is given by a time-dependent vector potential. We review from [BoGKS] the main tools that enter the derivation of the linear response, in particular the time evolution propagators. In Section 4 we compute rigorously the linear response of the system forced by a time dependent electric field. We do it along the lines of [BoGKS] but within the framework of the noncommutative L^p -spaces presented in Section 2. The derivation is achieved in several steps. First we set up the Liouville equation which describes the time evolution of the density matrix under the action of a time-dependent electric field (Theorem 4.1). In a standard way, this evolution equation can be written as an integral equation, the so-called Duhamel formula. Second, we compute the net current per unit volume induced by the electric field and prove that it is differentiable with respect to the electric field at zero field. This yields in fair generality the desired Kubo formula for the electric conductivity tensor, for any

non zero adiabatic parameter (Theorem 4.6 and Corollary 4.7). The adiabatic limit is then performed in order to compute the direct / ac conductivity at zero temperature (Theorem 4.8, Corollary 4.9 and Remark 4.11). In particular we recover the expected expression for the quantum Hall conductivity, the Kubo-Středa formula, as in [B, BES]. At positive temperature, we note that, while the existence of the electric conductivity *measure* can easily be derived from that Kubo formula [KM], proving that the conductivity itself, i.e. its density, exists and is finite remains out of reach.

Acknowledgement. We thank warmly Vladimir Georgescu for enlightening discussions on noncommutative integration, as well as A. Klein for useful comments.

2. COVARIANT MEASURABLE OPERATORS AND NONCOMMUTATIVE L^p -SPACES

In this section we construct the noncommutative L^p -spaces that are relevant for our analysis. We first recall the underlying Von Neumann algebra of observables and we equip it with the so called “trace per unit volume”. We refer to [D, Te] for the material. We shall skip some details and proofs for which we also refer to [Do].

2.1. Observables. Let \mathcal{H} be a separable Hilbert space (in our context $\mathcal{H} = L^2(\mathbb{R}^d)$). Let \mathcal{Z} be an abelian locally compact group and $U = \{U_a\}_{a \in \mathcal{Z}}$ a unitary projective representation of \mathcal{Z} on \mathcal{H} , i.e.

- $U_a U_b = \xi(a, b) U_{a+b}$, where $\xi(a, b) \in \mathbb{C}$, $|\xi(a, b)| = 1$;
- $U_e = 1$;
- $U_a^* = U_a^{-1} = \xi(a, -a)^{-1} U_{-a}$.

Now we take a set of orthogonal projections on \mathcal{H} , $\chi := \{\chi_a\}_{a \in \mathcal{Z}}$, $\mathcal{Z} \rightarrow \mathcal{B}(\mathcal{H})$. Such that if $a \neq b \Rightarrow \chi_a \chi_b = 0$ and $\sum_{a \in \mathcal{Z}} \chi_a = 1$. Furthermore one requires a covariance relation or a stability relation of χ under U i.e. $U_a \chi_b U_a^* = \chi_{a+b}$.

Next to this Hilbertian structure (representing the “physical” space), we consider a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ (representing the presence of the disorder) that is ergodic under the action of a group $\tau = \{\tau_a\}_{a \in \mathcal{Z}}$, that is,

- $\forall a \in \mathcal{Z}$, $\tau_a : \Omega \rightarrow \Omega$ is a measure preserving isomorphism;
- $\forall a, b \in \mathcal{Z}$, $\tau_a \circ \tau_b = \tau_{a+b}$;
- $\tau_e = 1$ where e is the neutral element of \mathcal{Z} and thus $\tau_a^{-1} = \tau_{-a}$, $\forall a \in \mathcal{Z}$;
- If $\mathcal{A} \in \mathcal{F}$ is invariant under τ , then $\mathbb{P}(\mathcal{A}) = 0$ or 1 .

With these two structures at hand we define the Hilbert space

$$\tilde{\mathcal{H}} = \int_{\Omega}^{\oplus} \mathcal{H} \, d\mathbb{P}(\omega) := L^2(\Omega, \mathbb{P}, \mathcal{H}) \simeq \mathcal{H} \otimes L^2(\Omega, \mathbb{P}), \quad (2.1)$$

equipped with the inner product

$$\langle \varphi, \psi \rangle_{\tilde{\mathcal{H}}} = \int_{\Omega} \langle \varphi(\omega), \psi(\omega) \rangle_{\mathcal{H}} \, d\mathbb{P}(\omega), \quad \forall \varphi, \psi \in \tilde{\mathcal{H}}^2. \quad (2.2)$$

We are interested in bounded operators on $\tilde{\mathcal{H}}$ that are decomposable elements $A = (A_{\omega})_{\omega \in \Omega}$, in the sense that they commute with the diagonal ones. Measurable operators are defined as decomposable operators such that for all measurable vector's field $\{\varphi(\omega), \omega \in \Omega\}$, the set $\{A(\omega)\varphi(\omega), \omega \in \Omega\}$ is measurable too. Measurable decomposable operators are called essentially bounded if $\omega \rightarrow \|A_{\omega}\|_{\mathcal{L}(\mathcal{H})}$ is a element of $L^{\infty}(\Omega, \mathbb{P})$. We set, for such A 's,

$$\|A\|_{\mathcal{L}(\tilde{\mathcal{H}})} = \|A\|_{\infty} = \text{ess - sup}_{\Omega} \|A(\omega)\|, \quad (2.3)$$

and define the following von Neumann algebra

$$\mathcal{K} = L^\infty(\Omega, \mathbb{P}, \mathcal{L}(\mathcal{H})) = \{A : \Omega \rightarrow \mathcal{L}(\mathcal{H}), \|A\|_\infty < \infty\}. \quad (2.4)$$

There exists an isometric isomorphism between \mathcal{K} and decomposable operators on $\mathcal{L}(\tilde{\mathcal{H}})$.

We shall work with observables of \mathcal{K} that satisfy the so-called covariant property.

Definition 2.1. *A $\in \mathcal{K}$ is covariant iff*

$$U_a A(\omega) U_a^* = A(\tau_a \omega), \quad \forall a \in \mathcal{Z}, \forall \omega \in \Omega. \quad (2.5)$$

We set

$$\mathcal{K}_\infty = \{A \in \mathcal{K}, A \text{ is covariant}\}. \quad (2.6)$$

If $\tilde{U}_a := U_a \otimes \tau(-a)$, and $\tilde{U} = (\tilde{U}_a)_{a \in \mathcal{Z}}$, we note that

$$\mathcal{K}_\infty = \{A \in \mathcal{K}, \forall a \in \mathcal{Z}, [A, \tilde{U}_a] = 0\} \quad (2.7)$$

$$= \mathcal{K} \cap (\tilde{U})', \quad (2.8)$$

so that \mathcal{K}_∞ is again a von Neumann algebra.

2.2. Noncommutative integration. The von Neumann algebra \mathcal{K}_∞ is equipped with the faithful, normal and semi-finite trace

$$\mathcal{T}(A) := \mathbb{E}\{\text{tr}(\chi_0 A(\omega) \chi_0)\}, \quad (2.9)$$

where “tr” denotes the usual trace on the Hilbert space \mathcal{H} . The trace \mathcal{T} is called the trace per unit volume, since by the Birkhoff Ergodic Theorem, whenever $\mathcal{T}(|A|) < \infty$, one has

$$\mathcal{T}(A) = \lim_{|\Lambda_L| \rightarrow \infty} \frac{1}{|\Lambda_L|} \text{tr}(\chi_{\Lambda_L} A \chi_{\Lambda_L}), \quad (2.10)$$

where $\Lambda_L \subset \mathcal{Z}$ and $\chi_{\Lambda_L} = \sum_{a \in \Lambda_L} \chi_a$. There is a natural topology associated to von Neumann algebras equipped with such a trace. It is defined by the following basis of neighborhoods:

$$N(\varepsilon, \delta) = \{A \in \mathcal{K}_\infty, \exists P \in \mathcal{K}_\infty^{proj}, \|AP\|_\infty < \varepsilon, \mathcal{T}(P^\perp) < \delta\}, \quad (2.11)$$

where $\mathcal{K}_\infty^{proj}$ denotes the set of projectors of \mathcal{K}_∞ . It is a well known fact that

$$A \in N(\varepsilon, \delta) \iff \mathcal{T}(\chi_{] \varepsilon, \infty[}(|A|)) \leq \delta. \quad (2.12)$$

As pointed out to us by V. Georgescu, this topology can also be generated by the following Frechet-norm on \mathcal{K}_∞ [Geo]:

$$\|A\|_{\mathcal{T}} = \inf_{P \in \mathcal{K}_\infty^{proj}} \max\{\|AP\|_\infty, \mathcal{T}(P^\perp)\}. \quad (2.13)$$

Let us denote by $\mathcal{M}(\mathcal{K}_\infty)$ the set of all \mathcal{T} -measurable operators, namely the completion of \mathcal{K}_∞ with respect to this topology. It is a well established fact from noncommutative integration that $\mathcal{M}(\mathcal{K}_\infty)$ is a Hausdorff topological $*$ -algebra.

Definition 2.2. *A linear subspace $\mathcal{E} \subseteq \mathcal{H}$ is called \mathcal{T} -dense if, $\forall \delta \in \mathbb{R}^+$, there exists a projection $P \in \mathcal{K}_\infty$ such that $P\mathcal{H} \subseteq \mathcal{E}$ and $\mathcal{T}(P^\perp) \leq \delta$.*

It turns out that any element A of $\mathcal{M}(\mathcal{K}_\infty)$ can be represented as an (possibly unbounded) operator, that we shall still denote by A , acting on $\tilde{\mathcal{H}}$ with a domain $D(A) = \{\varphi \in \tilde{\mathcal{H}}, A\varphi \in \tilde{\mathcal{H}}\}$ that is \mathcal{T} -densely defined. Then, adjoints, sums and products of elements of $\mathcal{M}(\mathcal{K}_\infty)$ are defined as usual adjoints, sums and products of unbounded operators.

For any $0 < p < \infty$, we set

$$L^p(\mathcal{K}_\infty) := \overline{\{x \in \mathcal{K}_\infty, \mathcal{T}(|x|^p) < \infty\}}^{\|\cdot\|_p} = \{x \in \mathcal{M}(\mathcal{K}_\infty), \mathcal{T}(|x|^p) < \infty\}, \quad (2.14)$$

where the second equality is actually a theorem. For $p \geq 1$, the spaces $L^p(\mathcal{K}_\infty)$ are Banach spaces in which $L^{p,o}(\mathcal{K}_\infty) := L^p(\mathcal{K}_\infty) \cap \mathcal{K}_\infty$ are dense by definition. For $p = \infty$, in analogy with the commutative case, we set $L^\infty(\mathcal{K}_\infty) = \mathcal{K}_\infty$.

Noncommutative Hölder inequalities hold: for any $0 < p, q, r \leq \infty$ so that $p^{-1} + q^{-1} = r^{-1}$, if $A \in L^p(\mathcal{K}_\infty)$ and $B \in L^q(\mathcal{K}_\infty)$, then the product $AB \in \mathcal{M}(\mathcal{K}_\infty)$ belongs to $L^r(\mathcal{K}_\infty)$ with

$$\|AB\|_r \leq \|A\|_p \|B\|_q. \quad (2.15)$$

In particular, for all $A \in L^\infty(\mathcal{K}_\infty)$ and $B \in L^p(\mathcal{K}_\infty)$,

$$\|AB\|_p \leq \|A\|_\infty \|B\|_p \text{ and } \|BA\|_p \leq \|A\|_\infty \|B\|_p, \quad (2.16)$$

so that $L^p(\mathcal{K}_\infty)$ -spaces are \mathcal{K}_∞ two-sided submodules of $\mathcal{M}(\mathcal{K}_\infty)$. As another consequence, bilinear forms $L^{p,o}(\mathcal{K}_\infty) \times L^{q,o}(\mathcal{K}_\infty) \ni (A, B) \mapsto \mathcal{T}(AB) \in \mathbb{C}$ continuously extends to bilinear maps defined on $L^p(\mathcal{K}_\infty) \times L^q(\mathcal{K}_\infty)$.

Lemma 2.3. *Let $A \in L^p(\mathcal{K}_\infty)$, $p \in [1, \infty[$ be given, and suppose $\mathcal{T}(AB) = 0$ for all $B \in L^q(\mathcal{K}_\infty)$, $p^{-1} + q^{-1} = 1$. Then $A = 0$.*

The case $p = 2$ is of particular interest since $L^2(\mathcal{K}_\infty)$ equipped with the sesquilinear form $\langle A, B \rangle_{L^2} = \mathcal{T}(A^*B)$ is a Hilbert space. The corresponding norm reads

$$\|A\|_2^2 = \int_\Omega \text{tr}(\chi_0 A_\omega^* A_\omega \chi_0) d\mathbb{P}(\omega) = \int_\Omega \|A_\omega \chi_0\|_2^2 d\mathbb{P}(\omega). \quad (2.17)$$

(Where $\|\cdot\|_2$ denotes the Hilbert-Schmidt norm.) From the case $p = 2$, we can derive the centrality of the trace. Indeed, by covariance and using the centrality of the usual trace, it follows that $\mathcal{T}(AB) = \mathcal{T}(BA)$ whenever $A, B \in \mathcal{K}_\infty$. By density we get the following lemma.

Lemma 2.4. *Let $A \in L^p(\mathcal{K}_\infty)$ and $B \in L^q(\mathcal{K}_\infty)$, $p^{-1} + q^{-1} = 1$ be given. Then $\mathcal{T}(AB) = \mathcal{T}(BA)$.*

We shall also make use of the following observation.

Lemma 2.5. *Let $A \in L^p(\mathcal{K}_\infty)$ and (B_n) a sequence of elements of \mathcal{K}_∞ that converges strongly to $B \in \mathcal{K}_\infty$. Then AB_n converges to AB in $L^p(\mathcal{K}_\infty)$.*

2.3. Commutators of measurable covariant operators. Let H be a decomposable (unbounded) operator affiliated to \mathcal{K}_∞ with domain \mathcal{D} , and $A \in \mathcal{M}(\mathcal{K}_\infty)$. In particular H need not be \mathcal{T} -measurable, i.e. in $\mathcal{M}(\mathcal{K}_\infty)$. If there exists a \mathcal{T} -dense domain \mathcal{D}' such that $A\mathcal{D}' \subset \mathcal{D}$, then HA is well defined, and if in addition the product is \mathcal{T} -measurable then we write $HA \in \mathcal{M}(\mathcal{K}_\infty)$. Similarly, if \mathcal{D} is \mathcal{T} -dense and the range of $H\mathcal{D} \subset \mathcal{D}(A)$, then AH is well defined, and if in addition the product is \mathcal{T} -measurable then we write $AH \in \mathcal{M}(\mathcal{K}_\infty)$.

Definition 2.6. *We define the following (generalized) commutators:*

(i): *If $A \in \mathcal{M}(\mathcal{K}_\infty)$ and $B \in \mathcal{K}_\infty$,*

$$[A, B] := AB - BA \in \mathcal{M}(\mathcal{K}_\infty), \quad [B, A] := -[A, B]. \quad (2.18)$$

(ii): *If $A \in L^p(\mathcal{K}_\infty)$, $B \in L^q(\mathcal{K}_\infty)$, $p, q \geq 1$ such that $p^{-1} + q^{-1} = 1$, then*

$$[A, B] := AB - BA \in L^1(\mathcal{K}_\infty). \quad (2.19)$$

(iii): Let H be affiliated to \mathcal{K}_∞ . If $A \in \mathcal{M}(\mathcal{K}_\infty)$ is such that HA and AH are in $\mathcal{M}(\mathcal{K}_\infty)$, then

$$[H, A] := HA - AH \in \mathcal{M}(\mathcal{K}_\infty). \quad (2.20)$$

We shall need the following observations.

Lemma 2.7. 1) For any $A \in L^p(\mathcal{K}_\infty), B \in L^q(\mathcal{K}_\infty), p, q \geq 1, p^{-1} + q^{-1} = 1$, and $C_\omega \in \mathcal{K}_\infty$, we have

$$\mathcal{T}\{[C, A]B\} = \mathcal{T}\{C[A, B]\}. \quad (2.21)$$

2) For any $A, B \in \mathcal{K}_\infty$ and $C \in L^1(\mathcal{K}_\infty)$, we have

$$\mathcal{T}\{A[B, C]\} = \mathcal{T}\{[A, B]C\}. \quad (2.22)$$

3) Let $p, q \geq 1$ be such that $p^{-1} + q^{-1} = 1$. For any $A \in L^p(\mathcal{K}_\infty)$, resp. $B \in L^q(\mathcal{K}_\infty)$, such that $[H, A] \in L^p(\mathcal{K}_\infty)$, resp. $[H, B] \in L^q(\mathcal{K}_\infty)$, we have

$$\mathcal{T}\{[H, A]B\} = -\mathcal{T}\{A[H, B]\}. \quad (2.23)$$

2.4. Differentiation. A $*$ -derivation ∂ is a $*$ -morphism defined on a dense subalgebra of \mathcal{K}_∞ and such that:

- $\partial(AB) = \partial(A)B + A\partial(B)$
- $\partial(A + \lambda B) = \partial(A) + \lambda\partial(B)$
- $\partial(A^*) = \partial(A)^*$
- $[\alpha_a, \partial] = 0$ in the sense that $\alpha_a \circ \partial(A) = \partial \circ \alpha_a(A) \forall a \in \mathcal{Z}, \forall A \in \mathcal{K}_\infty$.

If $\partial_1, \dots, \partial_d$ are $*$ -derivations we define a non-commutative gradient by $\nabla := (\partial_1, \dots, \partial_d)$, densely defined on \mathcal{K}_∞ . We define a non-commutative Sobolev space

$$\mathcal{W}^{1,p}(\mathcal{K}_\infty) := \{A \in L^p(\mathcal{K}_\infty), \nabla A \in L^p(\mathcal{K}_\infty)\}. \quad (2.24)$$

3. THE SETTING: SCHRÖDINGER OPERATORS AND DYNAMICS

In this section we describe our background operators and recall from [BoGKS] the main properties we shall need in order to establish the Kubo formula, but within the framework of noncommutative integration when relevant (i.e. in Subsection 3.2).

3.1. Magnetic Schrödinger operators and time-dependent operators.

Throughout this paper we shall consider Schrödinger operators of general form

$$H = H(\mathbf{A}, V) = (-i\nabla - \mathbf{A})^2 + V \quad \text{on } L^2(\mathbb{R}^d), \quad (3.1)$$

where the magnetic potential \mathbf{A} and the electric potential V satisfy the Leinfelder-Simader conditions:

- $\mathbf{A}(x) \in L^4_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d)$ with $\nabla \cdot \mathbf{A}(x) \in L^2_{\text{loc}}(\mathbb{R}^d)$.
- $V(x) = V_+(x) - V_-(x)$ with $V_\pm(x) \in L^2_{\text{loc}}(\mathbb{R}^d)$, $V_\pm(x) \geq 0$, and $V_-(x)$ relatively bounded with respect to Δ with relative bound < 1 , i.e., there are $0 \leq \alpha < 1$ and $\beta \geq 0$ such that

$$\|V_-\psi\| \leq \alpha\|\Delta\psi\| + \beta\|\psi\| \quad \text{for all } \psi \in \mathcal{D}(\Delta). \quad (3.2)$$

Leinfelder and Simader have shown that $H(\mathbf{A}, V)$ is essentially self-adjoint on $C_c^\infty(\mathbb{R}^d)$ [LS, Theorem 3]. It has been checked in [BoGKS] that under these hypotheses $H(\mathbf{A}, V)$ is bounded from below:

$$H(\mathbf{A}, V) \geq -\frac{\beta}{(1-\alpha)} =: -\gamma + 1, \quad \text{so that } H + \gamma \geq 1. \quad (3.3)$$

We denote by x_j the multiplication operator in $L^2(\mathbb{R}^d)$ by the j^{th} coordinate x_j , and $\mathbf{x} := (x_1, \dots, x_d)$. We want to implement the adiabatic switching of a time dependent spatially uniform electric field $\mathbf{E}_\eta(t) \cdot \mathbf{x} = e^{\eta t} \mathbf{E}(t) \cdot \mathbf{x}$ between time $t = -\infty$ and time $t = t_0$. Here $\eta > 0$ is the adiabatic parameter and we assume that

$$\int_{-\infty}^{t_0} e^{\eta t} |\mathbf{E}(t)| dt < \infty. \quad (3.4)$$

To do so we consider the time-dependent magnetic potential $\mathbf{A}(t) = \mathbf{A} + \mathbf{F}_\eta(t)$, with $\mathbf{F}_\eta(t) = \int_{-\infty}^t \mathbf{E}_\eta(s) ds$. In other terms, the dynamics is generated by the time-dependent magnetic operator

$$H(t) = (-i\nabla - \mathbf{A} - \mathbf{F}_\eta(t))^2 + V(x) = H(\mathbf{A}(t), V), \quad (3.5)$$

which is essentially self-adjoint on $C_c^\infty(\mathbb{R}^d)$ with domain $\mathcal{D} := \mathcal{D}(H) = \mathcal{D}(H(t))$ for all $t \in \mathbb{R}$. One has (see [BoGKS, Proposition 2.5])

$$H(t) = H - 2\mathbf{F}_\eta(t) \cdot \mathbf{D}(\mathbf{A}) + \mathbf{F}_\eta(t)^2 \text{ on } \mathcal{D}(H), \quad (3.6)$$

where $\mathbf{D} = \mathbf{D}(\mathbf{A})$ is the closure of $(-i\nabla - \mathbf{A})$ as an operator from $L^2(\mathbb{R}^d)$ to $L^2(\mathbb{R}^d; \mathbb{C}^d)$ with domain $C_c^\infty(\mathbb{R}^d)$. Each of its components $\mathbf{D}_j = \mathbf{D}_j(\mathbf{A}) = (-i\frac{\partial}{\partial x_j} - \mathbf{A}_j)$, $j = 1, \dots, d$, is essentially self-adjoint on $C_c^\infty(\mathbb{R}^d)$.

To see that such a family of operators generates the dynamics a quantum particle in the presence of the time dependent spatially uniform electric field $\mathbf{E}_\eta(t) \cdot \mathbf{x}$, consider the gauge transformation

$$[G(t)\psi](x) := e^{i\mathbf{F}_\eta(t) \cdot x} \psi(x), \quad (3.7)$$

so that

$$H(t) = G(t) [(-i\nabla - \mathbf{A})^2 + V] G(t)^*. \quad (3.8)$$

Then if $\psi(t)$ obeys Schrödinger equation

$$i\partial_t \psi(t) = H(t)\psi(t), \quad (3.9)$$

one has, *formally*,

$$i\partial_t G(t)^* \psi(t) = [(-i\nabla - \mathbf{A})^2 + V + \mathbf{E}_\eta(t) \cdot x] G(t)^* \psi(t). \quad (3.10)$$

To summarize the action of the gauge transformation we recall the

Lemma 3.1. [BoGKS, Lemma 2.6] *Let $G(t)$ be as in (3.7). Then*

$$G(t)\mathcal{D} = \mathcal{D}, \quad (3.11)$$

$$H(t) = G(t)HG(t)^*, \quad (3.12)$$

$$\mathbf{D}(\mathbf{A} + \mathbf{F}_\eta(t)) = \mathbf{D}(\mathbf{A}) - \mathbf{F}_\eta(t) = G(t)\mathbf{D}(\mathbf{A})G(t)^*. \quad (3.13)$$

Moreover, $i[H(t), x_j] = 2\mathbf{D}(\mathbf{A} + \mathbf{F}_\eta(t))$ as quadratic forms on $\mathcal{D} \cap \mathcal{D}(x_j)$, $j = 1, 2, \dots, d$.

The key observation is that the general theory of propagators with a time dependent generator [Y, Theorem XIV.3.1] applies to $H(t)$. It thus yields the existence

of a two parameters family $U(t, s)$ of unitary operators, jointly strongly continuous in t and s , that solves the Schrödinger equation.

$$U(t, r)U(r, s) = U(t, s) \quad (3.14)$$

$$U(t, t) = I \quad (3.15)$$

$$U(t, s)\mathcal{D} = \mathcal{D}, \quad (3.16)$$

$$i\partial_t U(t, s)\psi = H(t)U(t, s)\psi \text{ for all } \psi \in \mathcal{D}, \quad (3.17)$$

$$i\partial_s U(t, s)\psi = -U(t, s)H(s)\psi \text{ for all } \psi \in \mathcal{D}. \quad (3.18)$$

We refer to [BoGKS, Theorem 2.7] for other relevant properties.

To compute the linear response, we shall make use of the following ‘‘Duhamel formula’’. Let $U^{(0)}(t) = e^{-itH}$. For all $\psi \in \mathcal{D}$ and $t, s \in \mathbb{R}$ we have [BoGKS, Lemma 2.8]

$$U(t, s)\psi = U^{(0)}(t-s)\psi + i \int_s^t U^{(0)}(t-r)(2\mathbf{F}_\eta(r) \cdot \mathbf{D}(\mathbf{A}) - \mathbf{F}_\eta(r)^2)U(r, s)\psi \, dr. \quad (3.19)$$

Moreover,

$$\lim_{|\mathbf{E}| \rightarrow 0} U(t, s) = U^{(0)}(t-s) \text{ strongly}. \quad (3.20)$$

3.2. Adding the randomness. Let (Ω, \mathbb{P}) be a probability space equipped with an ergodic group $\{\tau_a; a \in \mathbb{Z}^d\}$ of measure preserving transformations. We study operator-valued maps $A: \Omega \ni \omega \mapsto A_\omega$.

Throughout the rest of this paper we shall use the material of Section 2 with $\mathcal{H} = L^2(\mathbb{R}^d)$ and $\mathcal{Z} = \mathbb{Z}^d$. The projective representation of \mathbb{Z}^d on \mathcal{H} is given by magnetic translations $(U(a)\psi)(x) = e^{ia \cdot Sx}\psi(x-a)$, S being a given $d \times d$ real matrix. The projection χ_a is the characteristic function of the unit cube of \mathbb{R}^d centered at $a \in \mathbb{Z}^d$.

In our context natural $*$ -derivations arise:

$$\partial_j A := i[x_j, A], \quad j = 1, \dots, d, \quad \nabla A = i[\mathbf{x}, A]. \quad (3.21)$$

We shall now recall the material from [BoGKS, Section 4.3]. Proofs of assertions are extensions of the arguments of [BoGKS] to the setting of $L^p(\mathcal{K}_\infty)$ -spaces. We refer to [Do] for details.

We state the technical assumptions on our Hamiltonian of reference H_ω .

Assumption 3.2. *The ergodic Hamiltonian $\omega \mapsto H_\omega$ is a measurable map from the probability space (Ω, \mathbb{P}) to self-adjoint operators on \mathcal{H} such that*

$$H_\omega = H(\mathbf{A}_\omega, V_\omega) = (-i\nabla - \mathbf{A}_\omega)^2 + V_\omega, \quad (3.22)$$

almost surely, where \mathbf{A}_ω (V_ω) are vector (scalar) potential valued random variables which satisfy the Leinfelder-Simader conditions (see Subsection 3.1) almost surely. It is furthermore assumed that H_ω is covariant as in (2.5). We denote by H the operator $(H_\omega)_{\omega \in \Omega}$ acting on $\tilde{\mathcal{H}}$.

As a consequence $\|f(H_\omega)\| \leq \|f\|_\infty$ and $f(H) \in \mathcal{K}_\infty$ for every bounded Borel function f on the real line. In particular H is affiliated to \mathcal{K}_∞ . For \mathbb{P} -a.e. ω , let $U_\omega(t, s)$ be the unitary propagator associated to H_ω and described in Subsection 3.1. Note that $(U_\omega(t, s))_{\omega \in \Omega} \in \mathcal{K}_\infty$ (measurability in ω follows by construction

of $U_\omega(t, s)$, see [BoGKS]). For $A \in \mathcal{M}(\mathcal{K}_\infty)$ decomposable, let

$$\mathcal{U}(t, s)(A) := \int_{\Omega}^{\oplus} U_\omega(t, s) A_\omega U_\omega(s, t) d\mathbb{P}(\omega). \quad (3.23)$$

Then $\mathcal{U}(t, s)$ extends a linear operator on $\mathcal{M}(\mathcal{K}_\infty)$, leaving invariant $\mathcal{M}(\mathcal{K}_\infty)$ and $L^p(\mathcal{K}_\infty)$, $p \in]0, \infty]$, with

$$\mathcal{U}(t, r)\mathcal{U}(r, s) = \mathcal{U}(t, s), \quad (3.24)$$

$$\mathcal{U}(t, t) = I, \quad (3.25)$$

$$\{\mathcal{U}(t, s)(A)\}^* = \mathcal{U}(t, s)(A^*). \quad (3.26)$$

Moreover, $\mathcal{U}(t, s)$ is to a unitary on $L^2(\mathcal{K}_\infty)$ and an isometry in $L^p(\mathcal{K}_\infty)$, $p \in]0, \infty]$. In addition, $\mathcal{U}(t, s)$ is jointly strongly continuous in t and s on $L^p(\mathcal{K}_\infty)$, $p \in]0, \infty]$.

Pick $p > 0$. Let $A \in L^p(\mathcal{K}_\infty)$ be such that $H(r_0)A$ and $AH(r_0)$ are in $L^p(\mathcal{K}_\infty)$ for some $r_0 \in [-\infty, \infty)$. Then both maps $r \mapsto \mathcal{U}(t, r)(A) \in L^p(\mathcal{K}_\infty)$ and $t \mapsto \mathcal{U}(t, r)(A) \in L^p(\mathcal{K}_\infty)$ are differentiable in $L^p(\mathcal{K}_\infty)$, with (recalling Definition 2.6)

$$i\partial_r \mathcal{U}(t, r)(A) = -\mathcal{U}(t, r)([H(r), A]). \quad (3.27)$$

$$i\partial_t \mathcal{U}(t, r)(A) = [H(t), \mathcal{U}(t, r)(A)]. \quad (3.28)$$

Moreover, for $t_0 < \infty$ given, there exists $C = C(t_0) < \infty$ such that for all $t, r \leq t_0$,

$$\|(H(t) + \gamma)\mathcal{U}(t, r)(A)\|_p \leq C\|(H(r) + \gamma)A\|_p, \quad (3.29)$$

$$\|[H(t), \mathcal{U}(t, r)(A)]\|_p \leq C(\|(H(r) + \gamma)A\|_p + \|A(H(r) + \gamma)\|_p). \quad (3.30)$$

We note that in order to apply the above formula and in particular (3.27) and (3.28), it is actually enough to check that $(H(r_0) + \gamma)A$ and $A(H(r_0) + \gamma)$ are in $L^p(\mathcal{K}_\infty)$.

Whenever we want to keep track of the dependence of $U_\omega(t, s)$ on the electric field $\mathbf{E} = \mathbf{E}_\eta(t)$, we shall write $U_\omega(\mathbf{E}, t, s)$. When $\mathbf{E} = 0$, note that

$$U_\omega(\mathbf{E} = 0, t, s) = U_\omega^{(0)}(t - s) := e^{-i(t-s)H_\omega}. \quad (3.31)$$

For $A \in \mathcal{M}(\mathcal{K}_\infty)$ decomposable, we let

$$\mathcal{U}^{(0)}(r)(A) := \int_{\Omega}^{\oplus} U_\omega^{(0)}(r) A_\omega U_\omega^{(0)}(-r) d\mathbb{P}(\omega). \quad (3.32)$$

We still denote by $\mathcal{U}^{(0)}(r)(A)$ its extension to $\mathcal{M}(\mathcal{K}_\infty)$.

Proposition 3.3. *Let $p \geq 1$ be given. $\mathcal{U}^{(0)}(t)$ is a one-parameter group of operators on $\mathcal{M}(\mathcal{K}_\infty)$, leaving $L^p(\mathcal{K}_\infty)$ invariant. $\mathcal{U}^{(0)}(t)$ is an isometry on $L^p(\mathcal{K}_\infty)$, and unitary if $p = 2$. It is strongly continuous on $L^p(\mathcal{K}_\infty)$. We further denote by \mathcal{L}_p the infinitesimal generator of $\mathcal{U}^{(0)}(t)$ in $L^p(\mathcal{K}_\infty)$:*

$$\mathcal{U}^{(0)}(t) = e^{-it\mathcal{L}_p} \quad \text{for all } t \in \mathbb{R}. \quad (3.33)$$

The operator \mathcal{L}_p is usually called the Liouvillian. Let

$$\mathcal{D}_p^{(0)} = \{A \in L^p(\mathcal{K}_\infty), HA, AH \in L^p(\mathcal{K}_\infty)\}. \quad (3.34)$$

Then $\mathcal{D}_p^{(0)}$ is an operator core for \mathcal{L}_p (note that \mathcal{L}_2 is essentially self-adjoint on $\mathcal{D}_2^{(0)}$), and

$$\mathcal{L}_p(A) = [H, A] \quad \text{for all } A \in \mathcal{D}_p^{(0)}. \quad (3.35)$$

Moreover, for every $B_\omega \in \mathcal{K}_\infty$ there exists a sequence $B_{n,\omega} \in \mathcal{D}_\infty^{(0)}$ such that $B_{n,\omega} \rightarrow B_\omega$ as a bounded and \mathbb{P} -a.e.-strong limit.

We finish this list of properties with the following lemma about the Gauge transformations in spaces of measurable operators. The map

$$\mathcal{G}(t)(A) = G(t)AG(t)^* , \quad (3.36)$$

with $G(t) = e^{i \int_{-\infty}^t \mathbf{E}_\eta(s) \cdot \mathbf{x}}$ as in (3.7), is an isometry on $L^p(\mathcal{K}_\infty)$, for $p \in]0, \infty]$.

Lemma 3.4. *For any $p \in]0, \infty]$, the map $\mathcal{G}(t)$ is strongly continuous on $L^p(\mathcal{K}_\infty)$, and*

$$\lim_{t \rightarrow -\infty} \mathcal{G}(t) = I \text{ strongly} \quad (3.37)$$

on $L^p(\mathcal{K}_\infty)$. Moreover, if $A \in \mathcal{W}^{1,p}(\mathcal{K}_\infty)$, then $\mathcal{G}(t)(A)$ is continuously differentiable in $L^p(\mathcal{K}_\infty)$ with

$$\partial_t \mathcal{G}(t)(A) = \mathbf{E}_\eta(t) \cdot \nabla(\mathcal{G}(t)(A)). \quad (3.38)$$

4. LINEAR RESPONSE THEORY AND KUBO FORMULA

4.1. Adiabatic switching of the electric field. We now fix an initial equilibrium state of the system, i.e., we specify a density matrix ζ_ω which is in equilibrium, so $[\mathbf{H}_\omega, \zeta_\omega] = 0$. For physical applications, we would generally take $\zeta_\omega = f(\mathbf{H}_\omega)$ with f the Fermi-Dirac distribution at inverse temperature $\beta \in (0, \infty]$ and *Fermi energy* $E_F \in \mathbb{R}$, i.e., $f(E) = \frac{1}{1+e^{\beta(E-E_F)}}$ if $\beta < \infty$ and $f(E) = \chi_{(-\infty, E_F]}(E)$ if $\beta = \infty$; explicitly

$$\zeta_\omega = \begin{cases} F_\omega^{(\beta, E_F)} & := \frac{1}{1+e^{\beta(\mathbf{H}_\omega - E_F)}}, \quad \beta < \infty, \\ P_\omega^{(E_F)} & := \chi_{(-\infty, E_F]}(\mathbf{H}_\omega), \quad \beta = \infty. \end{cases} \quad (4.1)$$

However we note that our analysis allows for fairly general functions f [BoGKS]. We set $\zeta = (\zeta_\omega)_{\omega \in \Omega} \in \mathcal{K}_\infty$ but shall also write ζ_ω instead of ζ . That f is the Fermi-Dirac distribution plays no role in the derivation of the linear response. However computing the Hall conductivity itself (once the linear response performed) we shall restrict our attention to the zero temperature case with the *Fermi projection* $P_\omega^{(E_F)}$.

The system is described by the ergodic time dependent Hamiltonian $H_\omega(t)$, as in (3.5). Assuming the system was in equilibrium at $t = -\infty$ with the density matrix $\varrho_\omega(-\infty) = \zeta_\omega$, the time dependent density matrix $\varrho_\omega(t)$ is the solution of the Cauchy problem for the Liouville equation. Since we shall solve the evolution equation in $L^p(\mathcal{K}_\infty)$, we work with $H(t) = (H_\omega(t))_{\omega \in \Omega}$, as in Assumption 3.2.

The electric field $\mathbf{E}_\eta(t) \cdot \mathbf{x} = e^{\eta t} \mathbf{E}(t) \cdot \mathbf{x}$ is switched on adiabatically between $t = -\infty$ and $t = t_0$ (typically $t_0 = 0$). Depending on which conductivity one is interested, one may consider different forms for $\mathbf{E}(t)$. In particular $\mathbf{E}(t) = \mathbf{E}$ leads to the direct conductivity, while $\mathbf{E}(t) = e^{i\nu t} \mathbf{E}$ leads to the AC-conductivity at frequency ν ¹. The first one is relevant for studying the Quantum Hall effect (see subsection 4.4), while the second enters the Mott's formula [KLP, KLM].

We write

$$\zeta(t) = G(t)\zeta G(t)^* = \mathcal{G}(t)(\zeta), \quad \text{i.e.,} \quad \zeta(t) = f(H(t)). \quad (4.2)$$

Theorem 4.1. *Let $\eta > 0$ and assume that $\int_{-\infty}^t e^{\eta r} |\mathbf{E}(r)| dr < \infty$ for all $t \in \mathbb{R}$. Let $p \in [1, \infty[$. Assume that $\zeta \in \mathcal{W}^{1,p}(\mathcal{K}_\infty)$ and that $H(\nabla\zeta), (\nabla\zeta)H \in L^p(\mathcal{K}_\infty)$. The*

¹The AC-conductivity may be better defined using the form (4.23) as argued in [KLM].

Cauchy problem

$$\begin{cases} i\partial_t \varrho(t) = [H(t), \varrho(t)] \\ \lim_{t \rightarrow -\infty} \varrho(t) = \zeta \end{cases}, \quad (4.3)$$

has a unique solution in $L^p(\mathcal{K}_\infty)$, that is given by

$$\varrho(t) = \lim_{s \rightarrow -\infty} \mathcal{U}(t, s) (\zeta) \quad (4.4)$$

$$= \lim_{s \rightarrow -\infty} \mathcal{U}(t, s) (\zeta(s)) \quad (4.5)$$

$$= \zeta(t) - \int_{-\infty}^t dr e^{\eta r} \mathcal{U}(t, r) (\mathbf{E}(r) \cdot \nabla \zeta(r)). \quad (4.6)$$

We also have

$$\varrho(t) = \mathcal{U}(t, s) (\varrho(s)), \quad \|\varrho(t)\|_p = \|\zeta\|_p, \quad (4.7)$$

for all t, s . Furthermore, $\varrho(t)$ is non-negative, and if ζ is a projection, then so is $\varrho(t)$ for all t .

Remark 4.2. If the initial state ζ is of the form (4.1), then the hypotheses of Theorem 4.1 hold for any $p > 0$, provided $\zeta_\omega = P_\omega^{(E_F)}$ that E_F lies in a region of localization. This is true for suitable $\mathbf{A}_\omega, V_\omega$ and E_F , by the methods of, for example, [CH, W, GK1, GK2, GK3, BoGK, AENSS, U, GrHK] and for the models studied therein as well as in [CH, GK3]. The bound $\mathbb{E} \|\mathbf{x} | \zeta_\omega \chi_0\|^2 < \infty$ or equivalently $\nabla \zeta \in L^2(\mathcal{K}_\infty)$ is actually sufficient for our applications. For $p = 1, 2$, we refer to [BoGKS, Proposition 4.2] and [BoGKS, Lemma 5.4] for the derivation of these hypotheses from known results.

Proof of Theorem 4.1. Let us first define

$$\varrho(t, s) := \mathcal{U}(t, s) (\zeta(s)). \quad (4.8)$$

We get, as operators in $\mathcal{M}(\mathcal{K}_\infty)$,

$$\begin{aligned} \partial_s \varrho(t, s) &= i\mathcal{U}(t, s) ([H(s), \zeta(s)]) + \mathcal{U}(t, s) (\mathbf{E}_\eta(s) \cdot \nabla \zeta(s)) \\ &= \mathcal{U}(t, s) (\mathbf{E}_\eta(s) \cdot \nabla \zeta(s)), \end{aligned} \quad (4.9)$$

where we used (3.27) and Lemma 3.4. As a consequence, with $\mathbf{E}_\eta(r) = e^{\eta r} \mathbf{E}(r)$,

$$\varrho(t, t) - \varrho(t, s) = \int_s^t dr e^{\eta r} \mathcal{U}(t, r) (\mathbf{E}(r) \cdot \nabla \zeta(r)). \quad (4.10)$$

Since $\|\mathcal{U}(t, r) (\mathbf{E}(r) \cdot \nabla (\zeta(r)))\|_p \leq c_d |\mathbf{E}(r)| \|\nabla \zeta\|_p < \infty$, the integral is absolutely convergent by hypothesis on $\mathbf{E}_\eta(t)$, and the limit as $s \rightarrow -\infty$ can be performed in $L^p(\mathcal{K}_\infty)$. It yields the equality between (4.5) and (4.6). Equality of (4.4) and (4.5) follows from Lemma 3.4 which gives

$$\zeta = \lim_{s \rightarrow -\infty} \zeta(s) \text{ in } L^p(\mathcal{K}_\infty). \quad (4.11)$$

Since $\mathcal{U}(t, s)$ are isometries on $L^p(\mathcal{K}_\infty)$, it follows from (4.4) that $\|\varrho(t)\|_p = \|\zeta\|_p$. We also get $\varrho(t) = \varrho(t)^*$. Moreover, (4.4) with the limit in $L^p(\mathcal{K}_\infty)$ implies that $\varrho(t)$ is nonnegative.

Furthermore, if $\zeta = \zeta^2$ then $\varrho(t)$ can be seen to be a projection as follows. Note that $\nabla \zeta = \zeta (\nabla \zeta) + (\nabla \zeta) \zeta$. Hölder inequalities imply that $\zeta \in \mathcal{W}^{1,r}(\mathcal{K}_\infty)$ for any

$0 < r \leq p$, and that the previous analysis actually applies in any $L^r(\mathcal{K}_\infty)$, $0 < r \leq p$. Denoting by $\lim^{(p)}$ the limit in $L^p(\mathcal{K}_\infty)$, we have, with $r = p/2$,

$$\begin{aligned} \varrho(t) &= \lim_{s \rightarrow -\infty}^{(r)} \mathcal{U}(t, s) (\zeta) = \lim_{s \rightarrow -\infty}^{(r)} \mathcal{U}(t, s) (\zeta) \mathcal{U}(t, s) (\zeta) \\ &= \left\{ \lim_{s \rightarrow -\infty}^{(p)} \mathcal{U}(t, s) (\zeta) \right\} \left\{ \lim_{s \rightarrow -\infty}^{(p)} \mathcal{U}(t, s) (\zeta) \right\} = \varrho(t)^2. \end{aligned} \quad (4.12)$$

To see that $\varrho(t)$ is a solution of (4.3) in $L^p(\mathcal{K}_\infty)$, we differentiate the expression (4.6) using (3.28) and Lemma 3.4. We get

$$i\partial_t \varrho(t) = - \int_{-\infty}^t dr e^{\eta r} [H(t), \mathcal{U}(t, r) (\mathbf{E}(r) \cdot \nabla \zeta(r))] \quad (4.13)$$

$$= - \left[H(t), \left\{ \int_{-\infty}^t dr e^{\eta r} \mathcal{U}(t, r) (\mathbf{E}(r) \cdot \nabla \zeta(r)) \right\} \right] \quad (4.14)$$

$$\begin{aligned} &= \left[H(t), \left\{ \zeta(t) - \int_{-\infty}^t dr e^{\eta r} \mathcal{U}(t, r) (\mathbf{E}(r) \cdot \nabla \zeta(r)) \right\} \right] \\ &= [H(t), \varrho(t)]. \end{aligned} \quad (4.15)$$

The integral in (4.13) converges since by (3.30) ,

$$\| [H(t), \mathcal{U}(t, r) (\mathbf{E}(r) \cdot \nabla \zeta(r))] \|_p \leq 2C \| (H + \gamma) (\mathbf{E}(r) \cdot \nabla \zeta) \|_p. \quad (4.16)$$

Then we justify going from (4.13) to (4.14) by inserting a resolvent $(H(t) + \gamma)^{-1}$ and making use of (3.29).

It remains to show that the solution of (4.3) is unique in $L^p(\mathcal{K}_\infty)$. It suffices to show that if $\nu(t)$ is a solution of (4.3) with $\zeta = 0$ then $\nu(t) = 0$ for all t . We define $\tilde{\nu}^{(s)}(t) = \mathcal{U}(s, t)(\nu(t))$ and proceed by duality. Since $p \geq 1$, with pick q s.t. $p^{-1} + q^{-1} = 1$. If $A \in \mathcal{D}_q^{(0)}$, we have, using Lemma 2.4,

$$\begin{aligned} i\partial_t \mathcal{T} \left\{ A \tilde{\nu}^{(s)}(t) \right\} &= i\partial_t \mathcal{T} \left\{ \mathcal{U}(t, s)(A) \nu(t) \right\} \\ &= \mathcal{T} \left\{ [H(t), \mathcal{U}(t, s)(A)] \nu(t) \right\} + \mathcal{T} \left\{ \mathcal{U}(t, s)(A) \mathcal{L}_q(t)(\nu(t)) \right\} \\ &= -\mathcal{T} \left\{ \mathcal{U}(t, s)(A) \mathcal{L}_q(t)(\nu(t)) \right\} + \mathcal{T} \left\{ \mathcal{U}(t, s)(A) \mathcal{L}_q(t)(\nu(t)) \right\} = 0. \end{aligned} \quad (4.17)$$

We conclude that for all t and $A \in \mathcal{D}_q^{(0)}$ we have

$$\mathcal{T} \left\{ A \tilde{\nu}^{(s)}(t) \right\} = \mathcal{T} \left\{ A \tilde{\nu}^{(s)}(s) \right\} = \mathcal{T} \left\{ A \nu(s) \right\}. \quad (4.18)$$

Thus $\tilde{\nu}^{(s)}(t) = \nu(s)$ by Lemma 2.3, that is, $\nu(t) = \mathcal{U}(t, s)(\nu(s))$. Since by hypothesis $\lim_{s \rightarrow -\infty} \nu(s) = 0$, we obtain that $\nu(t) = 0$ for all t . \square

4.2. The current and the conductivity. The velocity operator \mathbf{v} is defined as

$$\mathbf{v} = \mathbf{v}(\mathbf{A}) = 2\mathbf{D}(\mathbf{A}), \quad (4.19)$$

where $\mathbf{D} = \mathbf{D}(\mathbf{A})$ is defined below (3.6). Recall that $\mathbf{v} = 2(-i\nabla - \mathbf{A}) = i[H, \mathbf{x}]$ on $C_c^\infty(\mathbb{R}^d)$. We also set $\mathbf{D}(t) = \mathbf{D}(\mathbf{A} + \mathbf{F}_\eta(t))$ as in (3.13), and $\mathbf{v}(t) = 2\mathbf{D}(t)$.

From now on $\varrho(t)$ will denote the unique solution to (4.3), given explicitly in (4.6). If $H(t)\varrho(t) \in L^p(\mathcal{K}_\infty)$ then clearly $\mathbf{D}_j(t)\varrho(t)$ can be defined as well by

$$\mathbf{D}_j(t)\varrho(t) = (\mathbf{D}_j(t)(H(t) + \gamma)^{-1}) ((H(t) + \gamma)\varrho(t)), \quad (4.20)$$

since $\mathbf{D}_j(t)(H(t) + \gamma)^{-1} \in \mathcal{K}_\infty$, and thus $\mathbf{D}_j(t)\varrho(t) \in L^p(\mathcal{K}_\infty)$.

Definition 4.3. Starting with a system in equilibrium in state ζ , the net current (per unit volume), $\mathbf{J}(\eta, \mathbf{E}; \zeta, t_0) \in \mathbb{R}^d$, generated by switching on an electric field \mathbf{E} adiabatically at rate $\eta > 0$ between time $-\infty$ and time t_0 , is defined as

$$\mathbf{J}(\eta, \mathbf{E}; \zeta, t_0) = \mathcal{T}(\mathbf{v}(t_0)\varrho(t_0)) - \mathcal{T}(\mathbf{v}\zeta). \quad (4.21)$$

As it is well known, the current is null at equilibrium:

Lemma 4.4. One has $\mathcal{T}(\mathbf{D}_j\zeta) = 0$ for all $j = 1, \dots, d$, and thus $\mathcal{T}(\mathbf{v}\zeta) = 0$.

Throughout the rest of this paper, we shall assume that the electric field has the form

$$\mathbf{E}(t) = \mathcal{E}(t)\mathbf{E} \quad (4.22)$$

where $\mathbf{E} \in \mathbb{C}^d$ gives the intensity of the electric in each direction while $|\mathcal{E}(t)| = \mathcal{O}(1)$ modulates this intensity as time varies. As pointed out above, the two cases of particular interest are $\mathcal{E}(t) = 1$ and $\mathcal{E}(t) = e^{i\nu t}$. We may however, as in [KLM], use the more general form

$$\mathcal{E}(t) = \int_{\mathbb{R}} e^{i\nu t} \hat{\mathcal{E}}(\nu) d\nu, \quad (4.23)$$

for suitable $\hat{\mathcal{E}}(\nu)$ (see [KLM]).

It is useful to rewrite the current (4.21), using (4.6) and Lemma 4.4, as

$$\begin{aligned} \mathbf{J}(\eta, \mathbf{E}; \zeta, t_0) &= \mathcal{T}\{2\mathbf{D}(0)(\varrho(t_0) - \zeta(t_0))\} \\ &= -\mathcal{T}\left\{2\int_{-\infty}^{t_0} dr e^{\eta r} \mathbf{D}(0)\mathcal{U}(t_0, r)(\mathbf{E}(r) \cdot \nabla\zeta(r))\right\}. \\ &= -\mathcal{T}\left\{2\int_{-\infty}^{t_0} dr e^{\eta r} \mathcal{E}(r)\mathbf{D}(0)\mathcal{U}(t_0, r)(\mathbf{E} \cdot \nabla\zeta(r))\right\}. \end{aligned} \quad (4.24)$$

The conductivity tensor $\sigma(\eta; \zeta, t_0)$ is defined as the derivative of the function $\mathbf{J}(\eta, \mathbf{E}; \zeta, t_0): \mathbb{R}^d \rightarrow \mathbb{R}^d$ at $\mathbf{E} = 0$. Note that $\sigma(\eta; \zeta, t_0)$ is a $d \times d$ matrix $\{\sigma_{jk}(\eta; \zeta, t_0)\}$.

Definition 4.5. For $\eta > 0$ and $t_0 \in \mathbb{R}$, the conductivity tensor $\sigma(\eta; \zeta, t_0)$ is defined as

$$\sigma(\eta; \zeta, t_0) = \partial_{\mathbf{E}}(\mathbf{J}(\eta, \mathbf{E}; \zeta, t_0))|_{\mathbf{E}=0}, \quad (4.25)$$

if it exists. The conductivity tensor $\sigma(\zeta, t_0)$ is defined by

$$\sigma(\zeta, t_0) := \lim_{\eta \downarrow 0} \sigma(\eta; \zeta, t_0), \quad (4.26)$$

whenever the limit exists.

4.3. Computing the linear response: a Kubo formula for the conductivity. The next theorem gives a ‘‘Kubo formula’’ for the conductivity at positive adiabatic parameter.

Theorem 4.6. Let $\eta > 0$. Under the hypotheses of Theorem 4.1 for $p = 1$, the current $\mathbf{J}(\eta, \mathbf{E}; \zeta, t_0)$ is differentiable with respect to \mathbf{E} at $\mathbf{E} = 0$ and the derivative $\sigma(\eta; \zeta)$ is given by

$$\sigma_{jk}(\eta; \zeta, t_0) = -\mathcal{T}\left\{2\int_{-\infty}^{t_0} dr e^{\eta r} \mathcal{E}(r)\mathbf{D}_j\mathcal{U}^{(0)}(t_0 - r)(\partial_k(\zeta))\right\}. \quad (4.27)$$

The analogue of [BES, Eq. (41)] and [SB2, Theorem 1] then holds:

Corollary 4.7. *Assume that $\mathcal{E}(t) = e^{i\nu t}$, $\nu \in \mathbb{R}$, then the conductivity $\sigma_{jk}(\eta; \zeta)$ is given by*

$$\sigma_{jk}(\eta; \zeta, t_0) = -e^{(\eta+i\nu)t_0} \mathcal{T} \left\{ 2\mathbf{D}_j (i\mathcal{L}_1 + i\nu + \eta)^{-1} (\partial_k \zeta) \right\}, \quad (4.28)$$

Proof of Theorem 4.6. For clarity, in this proof we display the argument \mathbf{E} in all functions which depend on \mathbf{E} . From (4.24) and $\mathbf{J}_j(\eta, 0; \zeta, t_0) = 0$ (Lemma 4.4), we have

$$\sigma_{jk}(\eta; \zeta, t_0) = - \lim_{E \rightarrow 0} 2\mathcal{T} \left\{ \int_{-\infty}^{t_0} dr e^{\eta r} \mathcal{E}(r) \mathbf{D}_{\mathbf{E},j}(0) \mathcal{U}(\mathbf{E}, 0, r) (\partial_k \zeta(\mathbf{E}, r)) \right\}. \quad (4.29)$$

First understand we can interchange integration and the limit $\mathbf{E} \rightarrow 0$, and get

$$\sigma_{jk}(\eta; \zeta, t_0) = -2 \int_{-\infty}^{t_0} dr e^{\eta r} \mathcal{E}(r) \lim_{E \rightarrow 0} \mathcal{T} \left\{ \mathbf{D}_j(\mathbf{E}, 0) \mathcal{U}(\mathbf{E}, 0, r) (\partial_k \zeta(\mathbf{E}, r)) \right\}. \quad (4.30)$$

The latter can easily be seen by inserting a resolvent $(H(t) + \gamma)^{-1}$ and making use of (3.29), the fact that $H\nabla\zeta \in L^1(\mathcal{K}_\infty)$, the inequality $|\mathcal{T}(A)| \leq \mathcal{T}(|A|)$ and dominated convergence.

Next, we note that for any s we have

$$\lim_{E \rightarrow 0} \mathcal{G}(\mathbf{E}, s) = I \text{ strongly in } L^1(\mathcal{K}_\infty), \quad (4.31)$$

which can be proven by a argument similar to the one used to prove Lemma 3.4. Along the same lines, for $B \in \mathcal{K}_\infty$ we have

$$\lim_{E \rightarrow 0} \mathcal{G}(\mathbf{E}, s)(B_\omega) = B_\omega \text{ strongly in } \mathcal{H}, \text{ with } \|\mathcal{G}(\mathbf{E}, s)(B)\|_\infty = \|B\|_\infty. \quad (4.32)$$

Recalling that $\mathbf{D}_{j,\omega}(\mathbf{E}, 0) = \mathbf{D}_{j,\omega} - \mathbf{F}_j(0)$ and that $\|\partial_k \zeta(\mathbf{E}, r)\|_1 = \|\partial_k \zeta\|_1 < \infty$, using Lemma 2.5,

$$\begin{aligned} \lim_{E \rightarrow 0} \mathcal{T} \left\{ \mathbf{D}_j(\mathbf{E}, 0) \mathcal{U}(\mathbf{E}, 0, r) (\partial_k \zeta(\mathbf{E}, r)) \right\} &= \lim_{E \rightarrow 0} \mathcal{T} \left\{ \mathbf{D}_j U(\mathbf{E}, 0, r) (\partial_k \zeta) U(\mathbf{E}, r, 0) \right\} \\ &= \lim_{E \rightarrow 0} \mathcal{T} \left\{ \mathbf{D}_j U(\mathbf{E}, 0, r) (\partial_k \zeta) U^{(0)}(r) \right\}, \end{aligned} \quad (4.33)$$

where we have inserted (and removed) the resolvents $(H(\mathbf{E}, r) + \gamma)^{-1}$ and $(H + \gamma)^{-1}$.

To proceed it is convenient to introduce a cutoff so that we can deal with \mathbf{D}_j as if it were in \mathcal{K}_∞ . Thus we pick $f_n \in C_c^\infty(\mathbb{R})$, real valued, $|f_n| \leq 1$, $f_n = 1$ on $[-n, n]$, so that $f_n(H)$ converges strongly to 1. Using Lemma 2.5, we have

$$\begin{aligned} \mathcal{T} \left\{ \mathbf{D}_j U(\mathbf{E}, 0, r) (\partial_k \zeta) U^{(0)}(r) \right\} &= \lim_{n \rightarrow \infty} \mathcal{T} \left\{ f_n(H) \mathbf{D}_j U(\mathbf{E}, 0, r) (\partial_k \zeta) U^{(0)}(r) \right\} \\ &= \lim_{n \rightarrow \infty} \mathcal{T} \left\{ U(\mathbf{E}, 0, r) ((\partial_k \zeta)(H + \gamma)) U^{(0)}(r) (H + \gamma)^{-1} f_n(H) \mathbf{D}_j \right\} \\ &= \mathcal{T} \left\{ U(\mathbf{E}, 0, r) ((\partial_k \zeta)(H + \gamma)) \left(U^{(0)}(r) (H + \gamma)^{-1} \mathbf{D}_j \right) \right\}, \end{aligned} \quad (4.34)$$

where we used the centrality of the trace, the fact that $(H + \gamma)^{-1}$ commutes with $U^{(0)}$ and then that $(H + \gamma)^{-1} \mathbf{D}_j \in \mathcal{K}_\infty$ in order to remove to limit $n \rightarrow \infty$. Finally, combining (4.33) and (4.34), we get

$$\lim_{E \rightarrow 0} \mathcal{T} \left\{ \mathbf{D}_j(\mathbf{E}, 0) \mathcal{U}(\mathbf{E}, 0, r) (\partial_k \zeta(\mathbf{E}, r)) \right\} \quad (4.35)$$

$$\begin{aligned} &= \mathcal{T} \left\{ U^{(0)}(-r) ((\partial_k \zeta)(H + \gamma)) U^{(0)}(r) (H + \gamma)^{-1} \mathbf{D}_j \right\} \\ &= \mathcal{T} \left\{ \mathbf{D}_j U^{(0)}(-r) (\partial_k \zeta) \right\}. \end{aligned} \quad (4.36)$$

The Kubo formula (4.27) now follows from (4.30) and (4.36). \square

4.4. The Kubo-Středa formula for the Hall conductivity. Following [BES, AG], we now recover the well-known Kubo-Středa formula for the Hall conductivity at zero temperature (see however Remark 4.11 for AC-conductivity). To that aim we consider the case $\mathcal{E}(t) = 1$ and $t_0 = 0$. Recall Definition 4.5. We write

$$\sigma_{j,k}^{(E_f)} = \sigma_{j,k}(P^{(E_f)}, 0), \text{ and } \sigma_{j,k}^{(E_f)}(\eta) = \sigma_{j,k}(\eta; P^{(E_f)}, 0). \quad (4.37)$$

Theorem 4.8. *Take $\mathcal{E}(t) = 1$ and $t_0 = 0$. If $\zeta = P^{(E_f)}$ is a Fermi projection satisfying the hypotheses of Theorem 4.1 with $p = 2$, we have*

$$\sigma_{j,k}^{(E_f)} = -iT \left\{ P^{(E_f)} \left[\partial_j P^{(E_f)}, \partial_k P^{(E_f)} \right] \right\}, \quad (4.38)$$

for all $j, k = 1, 2, \dots, d$. As a consequence, the conductivity tensor is antisymmetric; in particular $\sigma_{j,j}^{(E_f)} = 0$ for $j = 1, 2, \dots, d$.

Clearly the direct conductivity vanishes, $\sigma_{j,j}^{(E_f)} = 0$. Note that, if the system is time-reversible the off diagonal elements are zero in the region of localization, as expected.

Corollary 4.9. *Under the assumptions of Theorem 4.8, if $\mathbf{A} = 0$ (no magnetic field), we have $\sigma_{j,k}^{(E_f)} = 0$ for all $j, k = 1, 2, \dots, d$.*

We have the crucial following lemma for computing the Kubo-Středa formula, which already appears in [BES] (and then in [AG]).

Lemma 4.10. *Let $P \in \mathcal{K}^\infty$ be a projection such that $\partial_k P \in L^p(\mathcal{K}^\infty)$, then as operators in $\mathcal{M}(\mathcal{K}^\infty)$ (and thus in $L^p(\mathcal{K}^\infty)$),*

$$\partial_k P = [P, [P, \partial_k P]]. \quad (4.39)$$

Proof. Note that $\partial_k P = \partial_k P^2 = P \partial_k P + (\partial_k P) P$ so that multiplying left and right both sides by P implies that $P(\partial_k P)P = 0$. We then have, in $L^p(\mathcal{K}^\infty)$,

$$\begin{aligned} \partial_k P &= P \partial_k P + (\partial_k P) P = P \partial_k P + (\partial_k P) P - 2P(\partial_k P)P \\ &= P(\partial_k P)(1 - P) + (1 - P)(\partial_k P)P \\ &= [P, [P, \partial_k P]]. \end{aligned}$$

\square

Remark that Lemma (4.10) heavily relies on the fact P is a projection. We shall apply it to the situation of zero temperature, i.e. when the initial density matrix is the orthogonal projection $P^{(E_f)}$. The argument would not go through at positive temperature.

Proof of Theorem 4.8. We again regularize the velocity $\mathbf{D}_{j,\omega}$ with a smooth function $f_n \in \mathcal{C}_c^\infty(\mathbb{R})$, $|f_n| \leq 1$, $f_n = 1$ on $[-n, n]$, but this time we also require that $f_n = 0$ outside $[-n-1, n+1]$, so that $f_n \chi_{[-n-1, n+1]} = f_n$. Thus $\mathbf{D}_j f_n(H) \in L^{p,o}(\mathcal{K}^\infty)$, $0 < p \leq \infty$. Moreover

$$f_n(H)(2\mathbf{D}_j)f_n(H) = f_n(H)P_n(2\mathbf{D}_j)P_n f_n(H) = -f_n(H)\partial_j(P_n H)f_n(H) \quad (4.40)$$

where $P_n = P_n^2 = \chi_{[-n-1, n+1]}(H)$ so that HP_n is a bounded operator. We have, using the centrality of the trace \mathcal{T} , that

$$\tilde{\sigma}_{jk}^{(E_F)}(r) := -\mathcal{T} \left\{ 2\mathbf{D}_{j,\omega} \mathcal{U}^{(0)}(-r) (\partial_k P^{(E_F)}) \right\} \quad (4.41)$$

$$= -\lim_{n \rightarrow \infty} \mathcal{T} \left\{ \mathcal{U}^{(0)}(r) (f_n(H) 2\mathbf{D}_{j,\omega} f_n(H)) \partial_k P^{(E_F)} \right\}. \quad (4.42)$$

Using Lemma 2.7 and applying Lemma 4.10 applied to $P = P^{(E_F)}$, it follows that

$$\mathcal{T} \left\{ \mathcal{U}^{(0)}(r) (f_n(H) 2\mathbf{D}_j f_n(H)) \partial_k P^{(E_F)} \right\} \quad (4.43)$$

$$\begin{aligned} &= \mathcal{T} \left\{ \mathcal{U}^{(0)}(r) (f_n(H) 2\mathbf{D}_j f_n(H)) \left[P^{(E_F)}, \left[P^{(E_F)}, \partial_k P^{(E_F)} \right] \right] \right\} \\ &= \mathcal{T} \left\{ \mathcal{U}^{(0)}(r) \left(\left[P^{(E_F)}, \left[P^{(E_F)}, f_n(H) 2\mathbf{D}_j f_n(H) \right] \right] \partial_k P^{(E_F)} \right) \right\}, \\ &= -\mathcal{T} \left\{ \mathcal{U}^{(0)}(r) \left(\left[P^{(E_F)}, f_n(H) \left[P^{(E_F)}, \partial_j (HP_n) \right] f_n(H) \right] \partial_k P^{(E_F)} \right) \right\}, \end{aligned}$$

where we used that $P^{(E_F)}$ commutes with $U^{(0)}$ and $f_n(H)$, and (4.40). Now, as elements in $\mathcal{M}(\mathcal{K}^\infty)$,

$$\left[P^{(E_F)}, \partial_j HP_n \right] = \left[HP_n, \partial_j P^{(E_F)} \right]. \quad (4.44)$$

Since $[H, \partial_j P^{(E_F)}]$ is well defined by hypothesis, $f_n(H) [HP_n, \partial_j P^{(E_F)}] f_n(H)$ converges in L^p to the latter as n goes to infinity. Combining (4.42), (4.43), and (4.44), we get after taking $n \rightarrow \infty$,

$$\tilde{\sigma}_{jk}^{(E_F)}(r) = -\mathcal{T} \left\{ \mathcal{U}^{(0)}(r) \left(\left[P^{(E_F)}, \left[H, \partial_j P^{(E_F)} \right] \right] \partial_k P^{(E_F)} \right) \right\}. \quad (4.45)$$

Next,

$$\left[P^{(E_F)}, \left[H, \partial_j P^{(E_F)} \right] \right] = \left[H, \left[P^{(E_F)}, \partial_j P^{(E_F)} \right] \right], \quad (4.46)$$

so that, recalling Proposition 3.3,

$$\begin{aligned} \tilde{\sigma}_{jk}^{(E_F)}(r) &= -\mathcal{T} \left\{ \mathcal{U}^{(0)}(r) \left(\left[H, \left[P^{(E_F)}, \partial_j P^{(E_F)} \right] \right] \partial_k P^{(E_F)} \right) \right\} \\ &= -\left\langle e^{-ir\mathcal{L}} \mathcal{L}_2 \left(\left[P^{(E_F)}, \partial_j P^{(E_F)} \right] \right), \partial_k P^{(E_F)} \right\rangle_{L^2}, \end{aligned} \quad (4.47)$$

where $\langle A, B \rangle_{L^2} = \mathcal{T}(A^*B)$. Combining (4.27), (4.41), and (4.47), we get

$$\sigma_{jk}^{(E_F)}(\eta) = -\left\langle i(\mathcal{L}_2 + i\eta)^{-1} \mathcal{L}_2 \left(\left[P^{(E_F)}, \partial_j P^{(E_F)} \right] \right), \partial_k P^{(E_F)} \right\rangle_{L^2}. \quad (4.48)$$

It follows from the spectral theorem (applied to \mathcal{L}_2) that

$$\lim_{\eta \rightarrow 0} (\mathcal{L}_2 + i\eta)^{-1} \mathcal{L}_2 = P_{(\text{Ker } \mathcal{L}_2)^\perp} \text{ strongly in } L^2(\mathcal{K}_\infty), \quad (4.49)$$

where $P_{(\text{Ker } \mathcal{L}_2)^\perp}$ is the orthogonal projection onto $(\text{Ker } \mathcal{L}_2)^\perp$. Moreover, as in [BoGKS] one can prove that

$$\left[P^{(E_F)}, \partial_j P^{(E_F)} \right] \in (\text{Ker } \mathcal{L}_2)^\perp. \quad (4.50)$$

Combining (4.48), (4.49), (4.50), and Lemma 2.7, we get

$$\sigma_{j,k}^{(E_F)} = i \left\langle \left[P^{(E_F)}, \partial_j P^{(E_F)} \right], \partial_k P^{(E_F)} \right\rangle_{L^2} = -i \mathcal{T} \left\{ P^{(E_F)} \left[\partial_j P^{(E_F)}, \partial_k P^{(E_F)} \right] \right\},$$

which is just (4.38). \square

Remark 4.11. *If one is interested in the AC-conductivity, then the proof above is valid up to (4.48). In particular, with $\mathcal{E}(t) = e^{i\nu t}$, one obtains*

$$\sigma_{jk}^{(E_F)}(\eta) = - \left\langle i(\mathcal{L}_2 + \nu + i\eta)^{-1} \mathcal{L}_2 \left(\left[P^{(E_F)}, \partial_j P^{(E_F)} \right] \right), \partial_k P^{(E_F)} \right\rangle_{L^2}. \quad (4.51)$$

The limit $\eta \rightarrow 0$ can still be performed as in [KLM, Corollary 3.4]. It is the main achievement of [KLM] to be able to investigate the behaviour of this limit as $\nu \rightarrow 0$ in connection with Mott's formula.

REFERENCES

- [AENSS] Aizenman, M., Elgart, A., Naboko, S., Schenker, J.H., Stolz, G.: Moment Analysis for Localization in Random Schrödinger Operators. 2003 Preprint, math-ph/0308023.
- [AG] Aizenman, M., Graf, G.M.: Localization bounds for an electron gas, J. Phys. A: Math. Gen. **31**, 6783-6806, (1998).
- [AvSS] Avron, J., Seiler, R., Simon, B.: Charge deficiency, charge transport and comparison of dimensions. Comm. Math. Phys. **159**, 399-422 (1994).
- [B] Bellissard, J.: Ordinary quantum Hall effect and noncommutative cohomology. In *Localization in disordered systems (Bad Schandau, 1986)*, pp. 61-74. Teubner-Texte Phys. **16**, Teubner, 1988.
- [BES] Bellissard, J., van Elst, A., Schulz-Baldes, H.: The non commutative geometry of the quantum Hall effect. J. Math. Phys. **35**, 5373-5451 (1994).
- [BH] Bellissard, J., Hislop, P.: Smoothness of correlations in the Anderson model at strong disorder. Ann. Henri Poincaré **8**, 1-26 (2007).
- [BoGK] Bouclet, J.M., Germinet, F., Klein, A.: Sub-exponential decay of operator kernels for functions of generalized Schrödinger operators. Proc. Amer. Math. Soc. **132**, 2703-2712 (2004).
- [BoGKS] Bouclet, J.M., Germinet, F., Klein, A., Schenker, J.: Linear response theory for magnetic Schrödinger operators in disordered media, J. Funct. Anal. **226**, 301-372 (2005)
- [CH] Combes, J.M., Hislop, P.D.: Landau Hamiltonians with random potentials: localization and the density of states. Commun. Math. Phys. **177**, 603-629 (1996).
- [CGH] Combes, J.M., Germinet, F., Hislop, P.D.: Conductivity and current-current correlation measure. In preparation.
- [CoJM] Cornean, H.D., Jensen, A., Moldoveanu, V.: The Landauer-Büttiker formula and resonant quantum transport. Mathematical physics of quantum mechanics, 45-53, Lecture Notes in Phys., 690, Springer, Berlin, 2006.
- [CoNP] Cornean, H.D., Nenciu, G., Pedersen, T.: The Faraday effect revisited: general theory. J. Math. Phys. **47** (2006), no. 1, 013511, 23 pp.
- [D] Dixmier, J.: Les algèbres d'opérateurs dans l'espace Hilbertien (algèbres de von Neumann), Gauthier-Villars 1969 and Gabay 1996.
- [Do] N. Dombrowski. PhD Thesis. In preparation.
- [ES] Elgart, A., Schlein, B.: Adiabatic charge transport and the Kubo formula for Landau Type Hamiltonians. Comm. Pure Appl. Math. **57**, 590-615 (2004).
- [FS] Fröhlich, J., Spencer, T.: Absence of diffusion with Anderson tight binding model for large disorder or low energy. Commun. Math. Phys. **88**, 151-184 (1983)
- [Geo] Georgescu, V.: Private communication.
- [GK1] Germinet, F., Klein, A.: Bootstrap Multiscale Analysis and Localization in Random Media. Commun. Math. Phys. **222**, 415-448 (2001).
- [GK2] Germinet, F., Klein, A.: Operator kernel estimates for functions of generalized Schrödinger operators. Proc. Amer. Math. Soc. **131**, 911-920 (2003).
- [GK3] Germinet, F., Klein, A.: Explicit finite volume criteria for localization in continuous random media and applications. Geom. Funct. Anal. **13**, 1201-1238 (2003).
- [GrHK] Ghribi, F., Hislop, P., Klopp, F.: Localization for Schrödinger operators with random vector potentials. Contemp. Math. To appear.
- [KLP] Kirsch, W., Lenoble, O., Pastur, L.: On the Mott formula for the ac conductivity and binary correlators in the strong localization regime of disordered systems. J. Phys. A **36**, 12157-12180 (2003).

- [KLM] Klein, A., Lenoble, O., Müller, P.: On Mott's formula for the AC-conductivity in the Anderson model. *Annals of Math.* To appear.
- [KM] Klein, A, Müller, P.: The conductivity measure for the Anderson model. In preparation.
- [Ku] Kunz, H.: The Quantum Hall Effect for Electrons in a Random Potential. *Commun. Math. Phys.* **112**, 121-145 (1987).
- [LS] H. Leinfelder, C.G. Simader, *Schrödinger operators with singular magnetic potentials*, *Math. Z.* 176, 1-19 (1981).
- [MD] Mott, N.F., Davies, E.A.: *Electronic processes in non-crystalline materials*. Oxford: Clarendon Press 1971.
- [NB] Nakamura, S., Bellissard, J.: Low Energy Bands do not Contribute to Quantum Hall Effect. *Commun. Math. Phys.* **131**, 283-305 (1990).
- [Na] Nakano, F.: Absence of transport in Anderson localization. *Rev. Math. Phys.* **14**, 375-407 (2002).
- [P] Pastur, L., Spectral properties of disordered systems in the one-body approximation. *Comm. Math. Phys.* **75**, 179-196 (1980).
- [PF] Pastur, L., Figotin, A.: *Spectra of Random and Almost-Periodic Operators*. Springer-Verlag, 1992.
- [SB1] Schulz-Baldes, H., Bellissard, J.: Anomalous transport: a mathematical framework. *Rev. Math. Phys.* **10**, 1-46 (1998).
- [SB2] Schulz-Baldes, H., Bellissard, J.: A Kinetic Theory for Quantum Transport in Aperiodic Media. *J. Statist. Phys.* **91**, 991-1026 (1998).
- [St] Středa, P.: Theory of quantised Hall conductivity in two dimensions. *J. Phys. C.* **15**, L717-L721 (1982).
- [Te] Terp, M.: L^p spaces associated with von Neumann algebras. Notes, Math. Institutue, Copenhagen university 1981.
- [ThKNN] Thouless, D. J., Kohmoto, K., Nightingale, M. P., den Nijs, M.: Quantized Hall conductance in a two-dimensional periodic potential. *Phys. Rev. Lett.* **49**, 405-408 (1982).
- [U] Ueki, N.: Wegner estimates and localization for Gaussian random potentials. *Publ. Res. Inst. Math. Sci.* 40, 29-90 (2004).
- [W] Wang, W.-M.: Microlocalization, percolation, and Anderson localization for the magnetic Schrödinger operator with a random potential. *J. Funct. Anal.* **146**, 1-26 (1997).
- [Y] Yosida, K.: *Functional Analysis, 6th edition*. Springer-Verlag, 1980.

(Dombrowski) UNIVERSITÉ DE CERGY-PONTOISE, CNRS UMR 8088, LABORATOIRE AGM, DÉPARTEMENT DE MATHÉMATIQUES, SITE DE SAINT-MARTIN, 2 AVENUE ADOLPHE CHAUVIN, F-95302 CERGY-PONTOISE, FRANCE

E-mail address: ndombro@math.u-cergy.fr

(Germinet) UNIVERSITÉ DE CERGY-PONTOISE, CNRS UMR 8088, LABORATOIRE AGM, DÉPARTEMENT DE MATHÉMATIQUES, SITE DE SAINT-MARTIN, 2 AVENUE ADOLPHE CHAUVIN, F-95302 CERGY-PONTOISE, FRANCE

E-mail address: germinet@math.u-cergy.fr